

University of California at San Diego – Department of Physics – Prof. John McGreevy

Physics 230 Quantum Phases of Matter, Spring 2024

Assignment 1

Due 11pm Thursday, April 11, 2024

1. Simple stabilizer codes.

I've mentioned the toric code as an important solvable example of topological order. We can solve it because it is an example of what is called a *stabilizer code*. This means that the system is made from a bunch of qubits and all the terms in the Hamiltonian are made of Pauli X s and Z s¹ and all commute with each other. This problem is a warmup problem to get used to this idea.

(a) Consider the Hamiltonian on two qubits

$$-H = X_1 X_2 + Z_1 Z_2.$$

Show that the terms commute and that the groundstate is

$$\frac{|00\rangle + |11\rangle}{\sqrt{2}}.$$

(b) Consider the (non-local) Hamiltonian on N qubits

$$H_{\text{GHZ}} = -X_1 \cdots X_N - \sum_{i=1}^{N-1} Z_i Z_{i+1}. \quad (1)$$

Show that all the terms commute. Show that the groundstate is (the ‘GHZ state’ or ‘cat state’)


$$\frac{|00\dots 0\rangle + |11\dots 1\rangle}{\sqrt{2}}.$$

¹A comment about notation: the notation $\sigma_\ell^x, \sigma_\ell^z$ is pretty terrible (at least for someone with deteriorating eyesight like me) because the crucial information (x or z) is hidden in the superscript. Much better is to write

$$\sigma_\ell^x \equiv X_\ell, \sigma_\ell^z \equiv Z_\ell.$$

Also, I use $|0\rangle, |1\rangle$ to denote the ± 1 eigenstates of Z , and $|\pm\rangle$ to denote the ± 1 eigenstates of X .

- (c) [bonus] Show that the following circuit U produces the GHZ state from the product state $|0\rangle^{\otimes N}$.



Let me explain the notation. Each horizontal line represents a qubit. H represents the ‘hadamard gate’ acting on one qubit by $H|\uparrow\rangle = |+\rangle$, $H|\downarrow\rangle = |-\rangle$, *i.e.*

$$H = |+\rangle\langle\uparrow| + |-\rangle\langle\downarrow|. \quad (2)$$

The vertical line segments represent the ‘control-X gate’ that acts on two qubits by

$$\text{CX} = |0\rangle\langle 0| \otimes \mathbb{1} + |1\rangle\langle 1| \otimes X. \quad (3)$$

(The first qubit is the one with the black dot, and the second is the one with the circled +.)

Notice that this circuit has depth N .

- (d) [bonus] What state does U produce from $|1\rangle_1 \otimes |0\rangle^{\otimes N-1}$?
- (e) [bonus] Find the result of feeding the Hamiltonian $-\sum_i Z_i$ (whose ground-state is the product state $|0\rangle^{\otimes N}$) through the circuit, *i.e.* what is

$$U \left(-\sum_i Z_i \right) U^\dagger ?$$

Hint: use the rules for the action of CX by conjugation given in lecture.


2. Gauge theory can emerge from a local Hilbert space.

The Hilbert space of a gauge theory is a funny thing: states related by a gauge transformation are physically equivalent. In particular, it is not a tensor product over independent local Hilbert spaces associated with regions of space. Because of this, there is much hand-wringing about defining entanglement in gauge theory. The following is helpful for thinking about this. It is a realization of \mathbb{Z}_2 lattice gauge theory, beginning from a model with no redundancy in its Hilbert space. In this avatar it is due to [Kitaev](#) and is called the *toric code*.

To define the Hilbert space, put a qubit on every link ℓ of a lattice, say the 2d square lattice, so that $\mathcal{H} = \otimes_{\ell} \mathcal{H}_{\ell}$. Let $\sigma_{\ell}^x, \sigma_{\ell}^z$ be the associated Pauli operators, and recall that $\{\sigma_{\ell}^x, \sigma_{\ell}^z\} = 0$. $\mathcal{H}_{\ell} = \text{span}\{|\sigma_{\ell}^z = 1\rangle, |\sigma_{\ell}^z = -1\rangle\}$ is a useful basis for the Hilbert space of a single link.

One term in the hamiltonian is associated with each site $j \rightarrow A_j \equiv \prod_{l \in j} \sigma_l^z$ and one with each plaquette $p \rightarrow B_p \equiv \prod_{l \in \partial p} \sigma_l^x$, as indicated in the figure at right.

$$\mathbf{H} = -\Gamma_e \sum_j A_j - \Gamma_m \sum_p B_p.$$



- (a) Show that all these terms commute with each other.
- (b) The previous result means we can diagonalize the Hamiltonian by diagonalizing one term at a time. Let's imagine that $\Gamma_e \gg \Gamma_m$ so we'll minimize the 'star' terms A_j first. Which states satisfy the 'star condition' $A_j = 1$? In the σ^x basis there is an extremely useful visualization: we say a link l of $\hat{\Gamma}$ is covered with a segment of string (an electric flux line) if $\sigma_l^z = -1$ (so the electric field on the link is $\mathbf{e}_l = 1$) and is not covered if $\sigma_l^z = +1$ (so the electric field on the link is $\mathbf{e}_l = 0$): $\text{---} \equiv (\sigma_l^z = -1)$. Draw all possible configurations incident on a single vertex j and characterize which ones satisfy $A_j = 1$.
- (c) [bonus] What is the effect of adding a term $\Delta \mathbf{H} = \sum_{\ell} g \sigma_{\ell}^x$? Convince yourself that in the limit $\Gamma_e \gg \Gamma_m$, for energies $E \ll \Gamma_e$, this is identical to \mathbb{Z}_2 lattice gauge theory, where $A_j = 1$ is a discrete version of the Gauss law constraint.
- (d) Set $g = 0$ again. In the subspace of solutions of the star condition, find the groundstate(s) of the plaquette term. First consider a simply-connected region of lattice, then consider periodic boundary conditions.

3. Groundstate degeneracy and 1-form symmetry algebra.

- (a) Suppose we have a system with Hamiltonian H with string operators W_C and $V_{\tilde{C}}$ supported on closed curves, and commuting with H , and satisfying $W^N = V^N = 1$.

In all parts of this problem you should make the assumption that the string operators are *deformable*: W_C acts in the same way as $W_{C+\partial p}$ on groundstates.

Suppose $[W_C, W_{C'}] = 0, [V_{\check{C}}, \check{C}']$ for all curves but

$$W_C V_{\check{C}} = \omega^{\#(C \cap \check{C})} V_{\check{C}} W_C$$

where $\omega \equiv e^{\frac{2\pi i}{N}}$ and $\#(C \cap \check{C})$ is the number of intersection points of the curves. How many groundstates does such a system have on the two-torus (that is, with periodic boundary conditions on both spatial directions)?

This is what happens in the \mathbb{Z}_N toric code.

- (b) Now suppose in a different system we have just one set of string operators W_C satisfying

$$W_C W_{C'} = \omega^{\#(C \cap C')} W_{C'} W_C,$$

with the same definitions as above. How many groundstates does this system have on the two-torus?

This is what happens in the Laughlin fractional quantum Hall state with filling fraction $\frac{1}{N}$.

- (c) [Bonus problem] Redo the previous problems for a genus g Riemann surface, *i.e.* the surface of a donut with g handles.

4. Simplicial homology and the toric code. [Bonus]

The toric code is a physical realization of *homology*, a construction that extracts topological invariants of topological spaces. This problem explains the relation.

- (a) The toric code we've discussed so far has qbits on the links $\ell \in \Delta_1(\mathbf{\Delta})$ of a graph $\mathbf{\Delta}$. But the definition of the Hamiltonian involves more information than just the links of the graph: we have to know which vertices v lie at the boundaries of each link ℓ , and we have to know which links are boundaries of which faces. The Hamiltonian has two kinds of terms: a 'plaquette' operator $B_p = \prod_{\ell \in \partial p} X_\ell$ associated with each 2-cell (plaquette) $p \in \Delta_2(\mathbf{\Delta})$, and 'star' operators, $A_s = \prod_{\ell \in \partial^{-1}(s)} Z_\ell$, associated with each 0-cell (site) $s \in \Delta_0(\mathbf{\Delta})$. Here I've introduced some notation that will be useful, please be patient: Δ_k denotes a collection of k -dimensional polyhedra which I'll call k -simplices or more accurately k -cells – k -dimensional objects making up the space. (It is important that each of these objects is topologically a k -ball.) This information constitutes (part of) a *simplicial complex*, which says how these parts are glued together:

$$\Delta_d \xrightarrow{\partial} \Delta_{d-1} \xrightarrow{\partial} \cdots \Delta_1 \xrightarrow{\partial} \Delta_0 \tag{4}$$

where ∂ is the (signed) boundary operator. For example, the boundary of a link is $\partial \ell = s_1 - s_0$, the difference of the vertices at its ends. The boundary

of a face $\partial p = \sum_{\ell \in \partial p} \ell$ is the (oriented) sum of the edges bounding it. By $\partial^{-1}(s)$ I mean the set of links which contain the site s in their boundary (with sign).

Think of this collection of objects as a triangulation (or more generally some chopping-up) of a smooth manifold X . Convince yourself that the sequence of maps (4) is a complex in the sense that $\partial^2 = 0 \pmod{\text{two}}$.

- (b) [not actually a question] This means that the simplicial complex defines a set of homology groups, which are topological invariants of X , in the following way. (It is homology and not cohomology because ∂ decreases the degree k). To define these groups, we should introduce one more gadget, which is a collection of vector spaces over some ring R (for the ordinary toric code, $R = \mathbb{Z}_2$)

$$\Omega_p(\Delta, R), \quad p = 0 \dots d \equiv \dim(X)$$

basis vectors for which are p -simplices:

$$\Omega_p(\Delta, R) = \text{span}_R\{\sigma \in \Delta_p\}$$

– that is, we associate a(n orthonormal) basis vector to each p -simplex (which I’ve just called σ), and these vector spaces are made by taking linear combinations of these spaces, with coefficients in R . Such a linear combination of p -simplices is called a p -chain. It’s important that we can *add* (and subtract) p -chains, $C + C' \in \Omega_p$. A p -chain with a negative coefficient can be regarded as having the opposite orientation. We’ll see below how better to interpret the coefficients.

The boundary operation on Δ_p induces one on Ω_p . A chain C satisfying $\partial C = 0$ is called a *cycle*, and is said to be *closed*.

So the p th homology is the group of equivalence classes of p -cycles, modulo boundaries of $p + 1$ cycles:

$$H_p(X, R) \equiv \frac{\ker(\partial : \Omega_p \rightarrow \Omega_{p-1}) \subset \Omega_p}{\text{Im}(\partial : \Omega_{p+1} \rightarrow \Omega_p) \subset \Omega_p}$$

This makes sense because $\partial^2 = 0$ – the image of $\partial : \Omega_{p+1} \rightarrow \Omega_p$ is a subset of $\ker(\partial : \Omega_p \rightarrow \Omega_{p-1})$. It’s a theorem that the dimensions of these groups are the same for different (faithful-enough) discretizations Δ of X . Furthermore, their dimensions (as vector spaces over R) $b_p(X)$ contain (much of²) the same information as the Betti numbers defined by de Rham cohomology. For more information and proofs, see the great book by Bott and Tu, *Differential forms in algebraic topology*.

²I don’t want to talk about torsion homology.

- (c) A state of the toric code on a cell-complex Δ can be written (for the hamiltonian described above, this is in the basis where Z_ℓ is diagonal) as an element of $\Omega_1(X, \mathbb{Z}_2)$,

$$|\Psi\rangle = \sum_C \Psi(C) |C\rangle$$

where C is an assignment of an element of \mathbb{Z}_2 in X (the eigenvalue of Z_ℓ). For the case of \mathbb{Z}_2 coefficients, $1 = -1 \pmod{2}$ and we don't care about the orientations of the cells. Show that the conditions for a state $\Psi(C)$ to be a groundstate of the toric code ($A_s |\Psi\rangle = |\Psi\rangle \forall s$ and $B_p |\Psi\rangle = |\Psi\rangle \forall p$) are exactly those defining an element of $H_1(X, \mathbb{Z}_2)$.

- (d) Consider putting a spin variable on the p -simplices of Δ . More generally, let's put an N -dimensional hilbert space $\mathcal{H}_N \equiv \text{span}\{|n\rangle, n = 1..N\}$ on each p -simplex, on which act the operators

$$\mathbf{X} \equiv \sum_{n=1}^N |n\rangle \langle n| \omega^n = \begin{pmatrix} 1 & 0 & 0 & \dots \\ 0 & \omega & 0 & \dots \\ 0 & 0 & \omega^2 & \dots \\ 0 & 0 & 0 & \ddots \end{pmatrix}, \quad \mathbf{Z} \equiv \sum_{n=1}^N |n\rangle \langle n+1| = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \vdots & \vdots & \vdots & \ddots \\ 1 & 0 & 0 & \dots \end{pmatrix}$$

where $\omega^N = 1$ is an n th root of unity. If you haven't already, check that they satisfy the clock-shift algebra: $\mathbf{Z}\mathbf{X} = \omega\mathbf{X}\mathbf{Z}$. For $N = 2$ these are Pauli matrices and $\omega = -1$.

Consider the Hamiltonian

$$\mathbf{H}_p = -J_{p-1} \sum_{s \in \Delta_{p-1}} A_s - J_{p+1} \sum_{\mu \in \Delta_{p+1}} B_\mu - g_p \sum_{\sigma \in \Delta_p} \mathbf{Z}_\sigma$$

with

$$A_s \equiv \prod_{\sigma \in \partial^{-1}(s) \subset \Delta_p} \mathbf{Z}_\sigma$$

$$B_\mu \equiv \prod_{\sigma \in \partial \mu} \mathbf{X}_\sigma .$$

This is a lattice version of p -form \mathbb{Z}_N gauge theory, at a particular, special point in its phase diagram (the RG fixed point for the deconfined phase).

Show that

$$0 = [A_s, A_{s'}] = [B_\mu, B_{\mu'}] = [A_s, B_\mu], \quad \forall s, s', \mu, \mu'$$

so that for $g_p = 0$ this is solvable.

- (e) Show that the groundstates of \mathbf{H}_p (with $g_p = 0$) are in one-to-one correspondence with elements of $H_p(\Delta, \mathbb{Z}_N)$.