

## Physics 217 Winter 2016 Assignment 2

Due 2pm Thursday, January 28, 2016

This homework begins with two simple problems to warm up your brain.

1. **Reminder.** Check from the ‘bridge’ equation  $F = -T \log Z$  that  $F = E - TS$  where  $E$  is the average energy and  $S = -\partial_T F$  is the entropy.
2. What is the relationship between the spin-spin correlation function  $\langle s_i s_j \rangle$  and the probability that  $s_i$  and  $s_j$  are pointing in the same direction? (Hint:  $(1 + s_i s_j)/2$  is a projector.)
3. **Real-space RG for the SAW.** Consider again the problem of a self-avoiding walk on the square lattice. Construct an RG scheme with zoom factor  $\lambda = 3$  (so that nine sites of the fine lattice are represented by one site of the coarse lattice). Find the RG map  $K'(K)$ , find its fixed points and estimate the critical exponent at the nontrivial fixed point. Is it closer to the numerical result than the  $\lambda = 2$  schemes discussed in lecture and by Creswick?

In the following problems we consider a closed (periodic) chain of  $N$  classical spins  $s_i = \pm 1$  with Hamiltonian

$$H = -J \sum_i s_i s_{i+1} - h \sum_i s_i + \text{const}, \quad s_{N+1} = s_1$$

The partition function is  $Z(\beta J, \beta h) = \sum_{\{s\}} e^{-\beta H}$ ; let’s measure  $J, h$  in units of temperature, *i.e.* set  $\beta = 1$ .

#### 4. Ising model in 1d by transfer matrix.

- (a) Show that the partition function can be written as

$$Z = \text{tr}_2 T^N$$

where  $T$  is the  $2 \times 2$  matrix

$$T = \begin{pmatrix} e^{J+h} & e^{-J} \\ e^{-J} & e^{J-h} \end{pmatrix}$$

(called the *transfer matrix*) and  $\text{tr}_2 M = M_{11} + M_{22}$  denotes trace in this two-dimensional space. Express  $Z$  in terms of the eigenvalues of  $T$  and find the free energy density  $f = -\frac{T}{N} \log Z$  in the thermodynamic ( $N \rightarrow \infty$ ) limit. Plot the free energy for  $h = 0$  as a function of  $x = e^{-4J}$  for  $0 \leq x \leq 1$ .

- (b) Find an expression for the correlation function

$$G(m) \equiv \langle s_i s_{m+i} \rangle - \langle s_i \rangle \langle s_{i+m} \rangle$$

using the transfer matrix. Show that as  $N \rightarrow \infty$ ,

$$G(m) \sim e^{-m/\xi}$$

where  $\xi = \frac{1}{\log\left(\frac{\lambda_1}{\lambda_2}\right)}$  where  $\lambda_1 > \lambda_2$  are eigenvalues of  $T$ . Note that  $\xi \rightarrow \infty$  when  $\lambda_1 \rightarrow \lambda_2$ . For what values of  $\beta, h, J$  does this happen?

## 5. Decimation of 1d Ising model in a field.

Now suppose that  $N$  is even.

- (a) Decimate the even sites:

$$\sum_{s_{\text{even}}} e^{-H(s)} \equiv \Delta e^{H_{\text{eff}}(s_{\text{odd}})}.$$

More explicitly, identify the terms in  $H(s)$  that depend on any one even site,  $H_2(s)$  and define its contribution to  $H_{\text{eff}}$  by

$$\sum_{s_2} e^{-H_2(s)} \equiv \Delta e^{-\Delta H_{\text{eff}}(s_1, s_3)}$$

Rewriting  $H_{\text{eff}}(s_{\text{odd}}) = -J' \sum s s - h' \sum s - \text{const}$  in the usual form, find  $J', h'$  and the constant in terms of the microscopic parameters  $J, h$ .

- (b) Let  $w \equiv \tanh \beta J, v \equiv \tanh \beta h$ . Plot some RG trajectories in the  $v, w$  plane.  
(c) Find all the fixed points and compute the exponents near each of the fixed points.  
(d) **I may add another section to this problem.**